DESY 97–167
hep-ph/9709207
September 1997

arXiv:hep-ph/9709207v1 1 Sep 1997

Semi-analytic predictions for W pair production at 500 GeV

J. Biebel and T. Riemann

Deutsches Elektronen-Synchrotron DESY, Institut für Hochenergiephysik
IfH Zeuthen, Platanenallee 6, D-15738 Zeuthen, Germany

Abstract

We give a short review on the semi-analytic approach to off-shell W pair production at 500 GeV and above. Numerical calculations are performed with the Fortran program `GENTLE` v. 2.0 and its upgrade. Recent extensions include:
(i) angular distribution for the `CC11` process with LLA QED corrections;
(ii) anomalous couplings.
As an application we study the sensitivity of total cross-section and forward-backward asymmetry to anomalous couplings. A noticeable sensitivity from the asymmetry appears only for a parity-violating anomalous coupling.

Emails: `biebel@ifh.de`, `riemann@ifh.de`

W pair production in e^+e^- annihilation at high energies has to be studied as off-shell production with at least four fermions in the final state ([1, 2, 3] and references therein). A convenient method for this is Monte Carlo simulation. In several respects, the semi-analytic approach of the Fortran program `GENTLE`[1] may prove useful:

- Tests and comparisons of different numerical programs [13, 14];
- Scan of W pair excitation curve [15, 16, 17];
- Search for anomalous triple boson couplings [18, 19].

The latter topic is better studied by multi-differential distributions/helicity analysis with Monte Carlo programs [18, 13]; but with a limited luminosity a single differential distribution, derived in a semi-analytical approach, may be a reasonable alternative.

The prospects of WW physics at a linear collider with center-of-mass energy of 500 GeV or more have been summarized in section 5.1 of [3]. Here, we study as an example the discriminative power of W pair production with respect to parity violating and conserving anomalous triple boson couplings. We assume $\sqrt{s} = 500$ GeV and an integrated luminosity $\mathcal{L} = 50$ fb^{-1}. This corresponds to about 80 000 semi-leptonic W pair decays. We make a simple ansatz. Let us explain it with only one anomalous coupling A. For this case, the cross-section may be written as follows:

$$\sigma_{theor}^{(1)} \;=\; \sigma_{SM} + A\sigma_1 + A^2\sigma_{11}. \tag{1}$$

The anomalous coupling appears in the cross-section at most bilinearly. After having calculated $\sigma_{SM}, \sigma_1, \sigma_{11}$ with `GENTLE` (or another program) within a given model and for fixed experimental conditions, one may confront an experimental data point with the predictions and get, in general, two possible solutions for A (or none). In a next step, one may consider two anomalous couplings A and B at once, etc.:

$$\sigma_{theor}^{(2)} \;=\; \sigma_{SM} + A\sigma_1 + A^2\sigma_{11} + B\sigma_2 + B^2\sigma_{22} + AB\sigma_{12}. \tag{2}$$

For our study of sensitivities, we use $\sigma_{theor}^{(2)}$ for the simulation of $\sigma_{meas} \pm \sqrt{\sigma_{meas}/(6\mathcal{L})}$, the assumed measured cross-section with 1σ deviations of the counting rates for the sum of all semi-leptonic production channels. For definiteness, we use for σ_{meas} the standard model prediction and apply no experimental cuts. The solutions of eq. (2) are ellipses in the plane spanned by the two anomalous couplings. Allowed pairs of coupling values are located in the area between the two limiting ellipses.

For the analysis, we use σ_F and σ_B, the forward and backward cross-sections. The forward (backward) cross-section is defined by the requirement that the angle between the momenta of the e^+ and the W^+ is less (more) than 90°. We chose these observables since they may be used to form the total cross-section $\sigma_{tot} = \sigma_F + \sigma_B$ (arizing from cross-section parts even in the production angle) and the forward backward asymmetry $A_{FB} = (\sigma_F - \sigma_B)/\sigma_{tot}$ (arising from odd cross-section parts).

---

[1] A detailed technical documentation of `GENTLE`, version 2.0, may be found in [4]. The approach is described in [5] (general features), [6, 7] (W pairs), [8, 9] (ZZ pairs), [10] (ZH production), [11, 12] (angular distributions, anomalous couplings).



Figure 1: 1σ-bounds at 500 GeV for $\mathcal{L} = 50 fb^{-1}$.

With the definition of the anomalous couplings x_γ and x_Z we follow equation (3) of [18] which is a re-parameterization of equation (1) of [18]. Additionally, we look at the triple boson coupling coupling z_Z defined by the effective Lagrangian

$$\mathcal{L}_Z \quad = \quad - \frac{e z_Z}{m_W^2} \partial_\alpha \hat{Z}_{\rho\sigma} \left(W^{+\alpha} \overleftrightarrow{\partial^\rho} W^{-\sigma} - W^{+\sigma} \overleftrightarrow{\partial^\rho} W^{-\alpha} \right) . \tag{3}$$

In figure 1, we show two solutions of eq. (2) for different anomalous couplings. In the left figure, x_γ and x_Z are allowed to differ from zero. These are $\mathcal{P}$ conserving couplings and the two rings derived from σ_F and σ_B overlap almost totally. In the second figure, x_Z and z_Z are varied. Since z_Z is $\mathcal{P}$ violating, the allowed ranges overlap much less since the forward-backward asymmetry is more sensitive to this coupling. One further notes that σ_B is more sensitive to anomalous couplings than σ_F although the relative statistical error of the latter is much smaller.

Acknowledgement

We would like to thank the convener of the Subgroup for Study of Electroweak Gauge Bosons, Thorsten Ohl, for numerous discussions, hints, and numerical comparisons with the Fortran program `WOPPER` [20, 21].

References

[1] G. Altarelli, T. Sjöstrand, and F. Zwirner (eds.), "Physics at LEP2", preprint CERN, CERN 96–01 (1996).

[2] W. Beenakker *et. al.*, "WW cross-sections and distributions", in [1], Vol. 1, p. 79, `hep-ph/9602351`.

[3] ECFA/DESY LC Physics Working Group, E. Accomando *et. al.*, "Physics with e^+e^- linear colliders", preprint DESY, DESY 97-100 (1997), `hep-ph/9705442,` subm. to Phys. Repts.

[4] D. Bardin, J. Biebel, D. Lehner, A. Leike, A. Olchevski, and T. Riemann, "`GENTLE/4fan v. 2.0`: A program for the semianalytic calculation of predictions for the process $e^+e^- \rightarrow 4f$", preprint DESY, DESY-96-233 (1996), `hep-ph/9612409,` to appear in *Comput. Phys. Commun.*

[5] D. Bardin, M. Bilenky, D. Lehner, A. Olchevski, and T. Riemann, "Semi-analytical approach to four-fermion production in e^+e^- annihilation", in *Proc. Zeuthen Workshop on Elementary Particle Theory – Physics at LEP200 and Beyond, Teupitz, Germany, April 10-15, 1994, Nucl. Phys. (Proc. Suppl.) 37B* (T. Riemann and J. Blümlein, eds.), pp. 148–157, Elsevier, Amsterdam, 1994.

[6] D. Bardin, A. Olshevsky, M. Bilenky, and T. Riemann, *Phys. Lett.* **B308** (1993) 403–410, E: ibid., B357 (1995) 725.

[7] D. Bardin and T. Riemann, *Nucl. Phys.* **B462** (1996) 3–28.

[8] D. Bardin, A. Leike, and T. Riemann, *Phys. Lett.* **B344** (1995) 383–390.

[9] D. Bardin, D. Lehner, and T. Riemann, *Nucl. Phys.* **B477** (1996) 27–58.

[10] D. Bardin, A. Leike, and T. Riemann, *Phys. Lett.* **B353** (1995) 513–518.

[11] A. Leike, "Semianalytic distributions in four fermion neutral current processes", in *Proc. Workshop on Perspectives for Electroweak Interactions in* e^+e^- *Collisions, Ringberg, Germany, Feb. 5 - 8, 1995* (B. A. Kniehl, ed.), pp. 121–130, World Scientific, River Edge, N. Y., 1995, `hep-ph/9504358`.

[12] J. Biebel and T. Riemann, in preparation.

[13] D. Bardin *et. al.*, "Event generators for WW physics", in [1], Vol. 2, p. 3.

[14] F. Boudjema *et. al.*, "Standard model processes", in [1], Vol. 1, p. 207, `hep-ph/9601224`.

[15] Z. Kunszt *et. al.*, "Determination of the mass of the W boson", in [1], Vol. 1, p.141, `hep-ph/9602352`.

[16] ALEPH Collaboration, R. Barate *et. al.*, "Measurement of the W pair cross-section in e^+e^- collisions at 172 GeV", preprint CERN, CERN-PPE-97-102 (1997).

[17] L3 Collaboration, M. Acciarri *et. al.*, “Measurement of W pair cross-sections in e^+e^- interactions at $\sqrt{s} = 172$ GeV and W decay branching fractions”, preprint CERN, CERN-PPE-97-067 (1997).

[18] G. Gounaris *et. al.*, “Triple gauge boson couplings”, in [1], Vol. 1, p. 525, `hep-ph/9601233`.

[19] L3 Collaboration, M. Acciarri *et. al.*, *Phys. Lett.* **B398** (1997) 223–238.

[20] H. Anlauf, J. Biebel, A. Himmler, P. Manakos, T. Mannel, and W. Schonau, *Comput. Phys. Commun.* **79** (1994) 487–502.

[21] H. Anlauf, P. Manakos, T. Ohl, and H. D. Dahmen, “WOPPER, version 1.5: A Monte Carlo event generator for $e^+e^- \to (W^+W^-) \to 4f + n\gamma$ at LEP-2 and beyond”, preprint TH Darmstadt, IKDA-96-15 (1996).